

# Nonconstant Error Variance in Generalized Propensity Score Model

Doyoung Kim, Chanmin Kim  
Department of Statistics, Sungkyunkwan University

## Preliminaries

### Annotation

- $\mathbf{X}$  : Covariate
- $T$  : Treatment

### Parametric Generalized Propensity Score Model

- $T_i | \mathbf{X}_i \sim N(\mathbf{X}_i^T \beta, \sigma^2)$
- Propensity score  $R_i = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(T_i - \mathbf{X}_i^T \beta)^2}{2\sigma^2}\right)$

### Caliper Metric Matching<sup>1)</sup>

$$m_{GPS}(e, w) = \operatorname{argmin}_{j: w_j \in [w - \delta, w + \delta]} \|(\lambda e^*(w_j, \mathbf{X}_j), (1 - \lambda)w_j^*) - (\lambda e^*, (1 - \lambda)w^*)\|$$

- $w_j^*$  : jth standardized treatment
- $e^*(w_j^*, \mathbf{X}_j)$  : Standardized propensity score with jth treatment & jth covariate
- $\delta$  : Caliper parameter
- $\lambda$  : Scale parameter between 0 & 1

### Covariate Balance (Global Measure)<sup>1)</sup>

$$\left| \sum_{i=1}^I \sum_{k=1}^{m_i} n_{ik} \mathbf{X}_{ik}^* W_{ik}^* \right| < \varepsilon_1$$

- $\mathbf{X}_{ik}^* = \mathbf{S}_X^{\frac{1}{2}} (\mathbf{X}_{ik} - \bar{\mathbf{X}}_{ik})$ ,  $W_{ik}^* = \mathbf{S}_W^{\frac{1}{2}} (W_{ik} - \bar{W}_{ik})$ ,  $\varepsilon_1$  is a pre-specified threshold (0.1)<sup>2)</sup>
- $I = \left\lfloor \frac{w^1 - w^0}{2\delta} + \frac{1}{2} \right\rfloor$  where  $[w^0, w^1]$  is range of treatment of interest
- $m_i$  : the number of units within the block  $[w_i - \delta, w_i + \delta]$   
where  $\{w_1 = w^0 + \delta, \dots, w_I = w^0 + (2I - 1)\delta\}$

1) Wu, X., Mealli, F., Kioumourtzoglou, M.-A., Dominici, F. & Braun, D. (2020), 'Matching on Generalized Propensity Scores with Continuous Exposures'

2) Zhu, Y., Coffman, D. L. & Ghosh, D. (2015), 'A Boosting Algorithm for Estimating Generalized Propensity Scores with Continuous Treatments', *Journal of causal inference*


## Motivation

### Existence of heteroskedasticity

- Variance of parametric GPS model depends on certain  $\mathbf{X}$
- Range of treatment the observation takes varies

e.g.

- $\mathbf{X}$  : Age, Race, Sex, Region ...  
 $T$  : Air pollution exposure  
 $Y$  : Mortality among 65+ years old



### Variance Function Model<sup>3)</sup>

$$T_i = f(\mathbf{X}_i, \beta) + g(\mu_i, z_i, \theta) \varepsilon_i, \quad \varepsilon_i \sim N(0, \sigma^2) \leftrightarrow T_i | \mathbf{X}_i \sim N(\mathbf{X}_i^T \beta, \sigma^2 g^2(\mu_i, z_i, \theta))$$

- $\mu_i : f(\mathbf{X}_i, \beta)$
- $z_i$  : Subset of  $\mathbf{X}_i$
- $\theta$  : Coefficient(s) of variance function

- Graphical methods for defining the form of variance function

e.g.

$$g(\mu_i, z_i, \theta) = 1 + \theta_1 z_i + \theta_2 z_i^2$$

if absolute residuals  $|r_i|$  &  $z_i$  shows quadratic form

3) Carroll, R. J., Ruppert, D., (1988), 'Transformation and Weighting in Regression', Chapman and Hall

## Model Specification & Proposal<sup>2)3)</sup>

### 1) Estimating $\theta$ & $\beta$

- Obtain LS estimator  $\hat{\beta}_{LS}, \hat{\beta}^{(0)} = \hat{\beta}_{LS}, k = 0$

- Minimize

$$\sum_{i=1}^N \left\{ r_i^2 \prod_{j=1}^N g(\mu_j(\hat{\beta}^{(k)}), z_j, \theta)^{\frac{1}{n}} \right\}^2$$

to obtain  $\hat{\theta}$

- Use the estimated weights  $\hat{w}_i = \frac{1}{g^2(\mu_i(\hat{\beta}^{(k)}), z_i, \hat{\theta})}$   
to obtain  $\hat{\beta}_{GLS}$

- Set  $k = k + 1$ , let  $\hat{\beta}^{(k)} = \hat{\beta}_{GLS}$  and return to (2)

### 2) Implement GPS matching approach

$$(1) \text{ Estimate GPS assuming } T_i | \mathbf{X}_i \sim N(\mathbf{X}_i^T \beta, \sigma^2 g^2(\mu_i, z_i, \theta))$$

$$\text{Propensity score } R_i = \frac{1}{\sqrt{2\pi\sigma^2 g^2(\mu_i, z_i, \theta)}} \exp\left(-\frac{(T_i - \mathbf{X}_i^T \beta)^2}{2\sigma^2 g^2(\mu_i, z_i, \theta)}\right)$$

- Select parameters  $(\lambda, \delta)$  minimizing covariate balance through grid search
- Match individuals based on one of proposed caliper metric matching methods

#### <Proposed Method 1>

$$m_{GPS}(e, w) = \operatorname{argmin}_{j: w_j \in [w - \delta, w + \delta]} \|(\alpha v_j^*, \beta w_j^*, (1 - \alpha - \beta)(w_j, \mathbf{X}_j)) - (\alpha v^*, \beta w^*, (1 - \alpha - \beta)e^*)\|$$

#### <Proposed Method 2>

$$d_j = |\sigma^2 g(\mu_j, z_j, \theta) - \sigma^2 g(\mu_i, z_i, \theta)|$$

$$m_{GPS}(e, w) = \operatorname{argmin}_{j: w_j \in [w - \delta, w + \delta]} \|(\lambda e^*(w_j, \mathbf{X}_j), (1 - \lambda)w_j^*) - (\lambda e^*, (1 - \lambda)w^*)\| \text{ for } d_j^* < \gamma$$

- Impute  $\hat{Y}_j(w) = Y_{\text{matched}}^{\text{obs}}$  for  $j = 1, \dots, N$  for all predetermined treatment levels  $w$ .

- With obtained  $\hat{\mu}(w) = \hat{E}\{\hat{Y}_j(w)\}$ , get a smoothed average causal treatment-response function

## Simulation

### Settings


- $n = 200$ , simulation replicates = 100
- Covariate :  $\mathbf{X}_1 - \mathbf{X}_4 \sim N(0, I_4)$ ,  $\mathbf{X}_5 \sim U(-2, 2)$ ,  $\mathbf{X}_6 \sim U(-3, 3)$
- Treatment :  $10 + 0.9\mathbf{X}_1 + 1.8\mathbf{X}_2 - 0.9\mathbf{X}_3 + 0.9\mathbf{X}_4 + 0.9\mathbf{X}_5 + 0.9\mathbf{X}_6$
- Outcome :  $Y | W, \mathbf{X} \sim N\{\mu(W, \mathbf{X}), 1.5^2\}$
- $\mu(W, \mathbf{X}) = 15 + 1.2\mathbf{X}_1 + 1.2\mathbf{X}_2 + 1.2\mathbf{X}_3 + 1.2\mathbf{X}_4 + 1.2\mathbf{X}_5 + \mathbf{X}_6 + 0.1W + 0.1W^2 + 0.25\mathbf{X}_2 W + 0.1\mathbf{X}_5 W$
- Variance function :  $\sigma^2 g^2(\mu_i, z_i, \theta) = 1^2(\theta_1 \mathbf{X}_4^2 + \theta_2 \mathbf{X}_4 + \theta_3 1.2\mathbf{X}_6)^2$   
 $(\theta_1, \theta_2, \theta_3) : (0.3, 1.2, 1.2), (0.4, 1.3, 1.3) \dots (1.0, 2.0, 2.0)$

### Metrics<sup>2)</sup>

- Absolute Bias :  $\frac{1}{M} \sum_{i=1}^M \left| \frac{1}{S} \sum_{s=1}^S \hat{Y}_s(w_i) - Y(w_i) \right|, \quad i = 1, \dots, M$
- MSE :  $\frac{1}{M} \sum_{i=1}^M \left[ \frac{1}{S} \sum_{s=1}^S \{\hat{Y}_s(w_i) - Y(w_i)\}^2 \right]^{1/2}$

### Results

- Covariate Balance



- Metrics

Default (0.3, 1.2, 1.2)			(0.4, 1.3, 1.3)			(0.5, 1.4, 1.4)		
	Absolute Bias	MSE		Absolute Bias	MSE		Absolute Bias	MSE
Prev	7.443	9.016	Prev	9.047	10.600	Prev	7.630	9.353
M1	7.518	8.637	M1	8.931	9.879	M1	7.602	9.064
M2	7.655	8.861	M2	9.403	10.706	M2	8.317	9.864

(0.6, 1.5, 1.5)			(0.7, 1.6, 1.6)			(0.8, 1.7, 1.7)		
	Absolute Bias	MSE		Absolute Bias	MSE		Absolute Bias	MSE
Prev	7.595	8.962	Prev	8.482	9.733	Prev	6.896	8.783
M1	7.416	8.390	M1	8.417	9.079	M1	6.812	7.932
M2	7.565	8.942	M2	8.845	9.949	M2	6.965	8.594

(1.0, 2.0, 2.0)		
	Absolute Bias	MSE
Prev	8.739	9.768
M1	8.880	9.520
M2	9.164	10.712

## Future Study

- Simulation metrics computation with marginal probabilities of treatment & full range of continuous treatment through kernel smoothing
- Application with heteroskedastic data